

# Independent origins of *Spiranthes* ×*kapnosperia* (Orchidaceae) and their nomenclatural implications

Matthew C. Pace<sup>1</sup>

<sup>1</sup> New York Botanical Garden, 2900 Southern Blvd., Bronx, New York, 10348, USA

Corresponding author: Matthew C. Pace ([mpace@nybg.org](mailto:mpace@nybg.org))

---

Academic editor: Timothée Le Péchon | Received 8 January 2023 | Accepted 28 April 2023 | Published 19 May 2023

---

**Citation:** Pace MC (2023) Independent origins of *Spiranthes* ×*kapnosperia* (Orchidaceae) and their nomenclatural implications. *PhytoKeys* 226: 89–100. <https://doi.org/10.3897/phytokeys.226.100062>

---

## Abstract

*Spiranthes* Rich. (Orchidaceae) is a commonly encountered but systematically and nomenclaturally challenging component of the North American orchid flora. Here, the evolutionary history and hybrid origin of the recently described *S. sheviakii* Hough and Young are critically examined. The available molecular data unambiguously support a hybrid origin of *S. cernua* (L.) Rich. × *S. ochroleuca* (Rydb.) Rydb. for *S. sheviakii*, the same parentage as the priority name *S. ×kapnosperia* M.C. Pace. As hybrid formulas can have only one correct name, *S. sheviakii* is a synonym of *S. ×kapnosperia*. It is likely that *S. ×kapnosperia* evolved independently at least twice in at least two widely disjunct locations.

## Keywords

hybrid speciation, Interior Lowlands, nomenclatural priority, species complex, *Spiranthes cernua*, *Spiranthes ochroleuca*

Species complexes continue to present some of the most impenetrable systematic challenges for evolutionary biology and conservation biology, and the challenges associated with their study are amplified when species within a complex hybridize (e.g., Arnold 2001; Fu et al. 2020), with challenging implications for nomenclature. Although Orchidaceae have been long seen as a model family for pre-zygotic barriers to hybridization, primarily due to documented or inferred pollinator specificity (Ackerman et al. 2023), a growing body of literature makes clear that reproductive barriers are often porous, and that hybridization plays an important role in the speciation of many orchid genera (e.g., *Dactylorhiza* Neck. ex Nevski (e.g., Pillon et al. 2007), *Epidendrum* (e.g., Pinheiro et al. 2010), *Ophrys* L. (e.g., Soliva and Widmer 2003), *Orchis* Tourn. ex L. (e.g., Jacquemyn




et al. 2012), *Platanthera* Rich. (e.g., Wettewa et al. 2020), and *Tolumnia* Raf. (e.g., Ackerman and Galarza-Pérez 1991)), making for ‘fuzzy’ species boundaries.

*Spiranthes* is one such orchid genus where renewed systematic attention has supported many previous hypotheses of hybridization (e.g., Dueck et al. 2014), in addition to the discovery of new hybrid taxa (e.g., Pace and Cameron 2017). Of the 44 currently accepted *Spiranthes* species (including nothospecies), 10 have molecular evidence to support a hybrid origin (Sun 1996; Arft and Ranker 1998; Szalanski et al. 2001, Dueck et al. 2014; Pace 2015, 2021; Pace and Cameron 2017, 2019; Surveswaran et al. 2018; Pace et al. 2019). These hybrid species do not only occur within complexes of closely related species (e.g., *S. ×stellata* P.M.Br., Dueck and K.M.Cameron), but between clades of species complexes that are often distantly related (e.g., *S. diluvialis* Sheviak). The *S. cernua* (L.) Rich. species complex has traditionally been regarded as systematically “intractable” (Sheviak 1982, 1991; Sheviak and Brown 2002), primarily due to the frequency of hybridization and cryptic speciation (Pace and Cameron 2017) and the variability of all taxa involved. Within the *S. cernua* species complex, the identity of *S. ochroleuca* (Rydb.) Rydb. has contributed significantly to systematic and nomenclatural challenges. For example, primarily due to the nature of the *Gyrostachys ochroleuca* Rydb. holotype (*Mrs Long s.n.*, drawing, NY barcode 9463, Fig. 1), and morphological similarities to other members of the complex, *S. ochroleuca* has either been treated as a synonym or variety of *S. cernua* for much of the last 90 years (e.g., Gleason and Cronquist 1963). It was only after the detailed work of Sheviak and Catling (1980), that *S. ochroleuca* was widely accepted as a species fully distinct from *S. cernua* (e.g., Pace and Freudenstein 2018). Despite this distinction, *S. cernua* s.s. and *S. ochroleuca* were still hypothesized to engage in frequent and widespread hybridization and introgression (Sheviak 1982; Sheviak and Brown 2002). Pace and Cameron (2017) presented the first molecular evidence for hybridization between *S. ochroleuca* and *S. cernua* s.s. in the southern Appalachians, describing this hybrid taxon as *S. ×kapnosperia* M.C. Pace.

The name *S. sheviakii* Hough and Young (2021) was recently described as a species of hybrid origin distributed from central New York to the greater Ohio River Valley, but Hough and Young (2021) were unusually vague about the parentage of *S. sheviakii*, writing “[this is] apparently the result of hybridization of *S. ochroleuca* with another member of the *S. cernua* species complex” (pg. 47). They included comparisons to *S. cernua*, *S. ochroleuca*, and *S. ×kapnosperia* in the diagnosis and throughout the discussion, noting that *S. sheviakii* is “intermediate in form” between *S. cernua* and *S. ochroleuca* (pg. 37), but did not give a full parentage to their newly proposed species. Curiously, Hough originally identified the type specimens of *S. sheviakii* as “*S. ×kapnosperia*, *S. cernua* × *S. ochroleuca*” (Fig. 1), indicating he was aware of its full parentage, or that he thought these plants were morphologically similar to *S. ×kapnosperia*.

After reviewing the relevant type specimens (Fig. 1) and the publicly available molecular data presented in Pace and Cameron (2017) and Hough and Young (2021), it is clear that both *S. ×kapnosperia* and *S. sheviakii* share a hybrid ancestry of *S. cernua* × *S. ochroleuca*, although the genetic patterns are differently expressed in the resulting regional hybrids. Appalachian *S. ×kapnosperia* displays a discordance between nuclear and chloroplast datasets: the chloroplast data (including *ndhF*)







indicates a maternal parent of *S. ochroleuca*, whereas the nuclear data (*ACO* and *nrITS*) indicate a paternal parentage of *S. cernua* (Table 2; Pace and Cameron 2017). The *ACO* dataset for Appalachian *S. ×kapnosperia* lacks major nucleotide ambiguities at points of differentiation between the parental species, sharing all of the unique molecular synapomorphies of *S. cernua* vs. *S. ochroleuca*. *Spiranthes sheviakii* also displays a discordance between nuclear and chloroplast data, although the discordance is slightly different than in Appalachian *S. ×kapnosperia*. The available *ndhJ* data for *S. sheviakii* clearly indicate a maternal parentage of *S. cernua*, as the samples share all of the same nucleotide patterns as *S. cernua* and are clearly different from *S. ochroleuca* (or any other member of the *S. cernua* species complex). However, *S. sheviakii* displays *ACO* nucleotide ambiguities at all of the exact and unique points of molecular differentiation between *S. cernua* and *S. ochroleuca* (Fig. 2). These ambiguities in the *ACO* dataset indicate a hybrid origin between *S. cernua* and *S. ochroleuca* for

**Table 1.** Comparison of *Spiranthes cernua*, *S. ×kapnosperia*, and *S. ochroleuca*.

Taxon	Distribution	Flower color	Flower position	Labellum color (Adaxial / Abaxial)	Abaxial gland shape	Lateral sepal position
<i>S. cernua</i>	Maritime Canada s. to n. FL, west to central PA, the interior lowland plateaus, and e. Texas	White	Perpendicular to stem to strongly nodding	White to very pale yellow / white	Conical, reduced	Sepal base held in-line with the profile of the flower, sepals ascending
<i>S. ×kapnosperia</i>	Southern Appalachian Mountains of NC, SC, and TN; southern Great Lakes Basin from central NY to IL, s. to Ohio River Valley	White to ivory	Nodding to slightly ascending	Pale yellow / pale yellow	Rounded	Sepal base held just below the profile of the flower, sepals ascending
<i>S. ochroleuca</i>	Maritime Canada s. to w. NC, west through the Southern Great Lakes Basin, disjunct in s. IL, IN, central KY, and TN	Ivory to ochroleucous	Slightly to strongly ascending	Deep yellow / deep yellow	Rounded (sometimes reddish)	Sepal base held below the profile of the flower, held low against the profile of the flower, downwardly falcate to ascending

**Table 2.** Inferred genetic contributions for *S. ×kapnosperia* sensu nov. Chloroplast regions include *matK*, *ndhJ*, *trnL-F*, *trnS-M*, and *ycf1* 3' (Pace and Cameron 2017). The only chloroplast region sampled in Hough and Young (2021) for a priori *S. sheviakii* is *ndhJ*.

Hybrid taxa	ACO (nuclear)	nrITS	Chloroplast
<i>S. ×kapnosperia</i> (Appalachian <i>S. ×kapnosperia</i> )	<i>S. cernua</i> s.s.	<i>S. cernua</i> s.s.	<i>S. ochroleuca</i>
<i>S. sheviakii</i> (Interior Lowland <i>S. ×kapnosperia</i> )	<i>S. cernua</i> s.s. + <i>S. ochroleuca</i>	[not sampled]	<i>S. cernua</i> s.s.



A.

105. S.magnicamporum_sm7h_NM	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
106. S.magnicamporum_sm8a_VA	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
107. S.magnicamporum_15f_TX	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
108. S.magnicamporum_15f_WI	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
109. S.magnicamporum_sm11a_VA	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
110. S.magnicamporum_sm15a_Ontario	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
111. S.magnicamporum_sm12a_GA	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
112. S.magnicamporum_sm13a_Ontario	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
113. S.magnicamporum_sm17a_TN	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
114. S.magnicamporum_sm19a_IL	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
115. S.magnicamporum_sm22c_WI	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
116. S.magnicamporum_sm25g_IL	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
117. S.magnicamporum_sm26k_IL	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
118. S.magnicamporum_sm28h_IN	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
119. S.magnicamporum_sm23e_IL	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
120. S.incurva_4v_OH	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
121. S.incurva_4t_WI	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
122. S.incurva_sm1c_WI	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
123. S.incurva_sm21r_WI	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
124. S.incurva_sm23b_IL	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
125. S.incurva_sc33a_Ontario	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
126. S.incurva_soch10a_VT	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
127. S.cernua_4cc_SC	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
128. S.cernua_4dd_NC	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
129. S.cernua_4l_TX	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
130. S.cernua_4m_FL	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
131. S.cernua_sc1b_DE	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
132. S.cernua_AR29_AR	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
133. S.cernua_sc9a_VA	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
134. S.cernua_sc15b_GA	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
135. S.×kapsheviakii_s20	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
136. S.×kapsheviakii_s26	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
137. S.×kapsheviakii_s33	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
138. S.×kapsheviakii_s36	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
139. S.×kapsheviakii_s38	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
140. S.×kapsheviakii_s40	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
141. S.×kapsnosperia_sc20a_NC	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
142. S.×kapsnosperia_sh2_NC	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
143. S.ochroleuca_4n_NY	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
144. S.ochroleuca_4wx_OH	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
145. S.ochroleuca_16b_VA	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
146. S.ochroleuca_16c_NH	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
147. S.ochroleuca_16f_NH	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
148. S.ochroleuca_16g_MI	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
149. S.ochroleuca_16h_NY	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
150. S.arcisepala_4u_OH	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
151. S.arcisepala_sc30a_OH	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
152. S.arcisepala_4y_NY	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
153. S.arcisepala_4z_OH	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT
154. S.arcisepala_NY1_NY	CCCATTTGTTATCACCTGTAGGATTAAAGTTGTTAAAGTTTTCATCGTCATCAAACTCTCTAAC	GTT	AAGC	TTTTTTTT

B.

105. S.magnicamporum_sm7h_NM	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
106. S.magnicamporum_sm8a_VA	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
107. S.magnicamporum_15f_TX	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
108. S.magnicamporum_15f_WI	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
109. S.magnicamporum_sm11a_VA	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
110. S.magnicamporum_sm15a_Ontario	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
111. S.magnicamporum_sm12a_GA	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
112. S.magnicamporum_sm13a_Ontario	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
113. S.magnicamporum_sm17a_TN	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
114. S.magnicamporum_sm19a_IL	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
115. S.magnicamporum_sm22c_WI	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
116. S.magnicamporum_sm25g_IL	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
117. S.magnicamporum_sm26k_IL	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
118. S.magnicamporum_sm28h_IN	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
119. S.magnicamporum_sm23e_IL	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
120. S.incurva_4v_OH	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
121. S.incurva_4t_WI	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
122. S.incurva_sm1c_WI	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
123. S.incurva_sm21r_WI	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
124. S.incurva_sm23b_IL	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
125. S.incurva_sc33a_Ontario	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
126. S.incurva_soch10a_VT	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
127. S.cernua_4cc_SC	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
128. S.cernua_4dd_NC	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
129. S.cernua_4l_TX	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
130. S.cernua_4m_FL	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
131. S.cernua_sc1b_DE	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
132. S.cernua_AR29_AR	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
133. S.cernua_sc9a_VA	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
134. S.cernua_sc15b_GA	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
135. S.×kapsheviakii_s20	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
136. S.×kapsheviakii_s26	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
137. S.×kapsheviakii_s33	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
138. S.×kapsheviakii_s36	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
139. S.×kapsheviakii_s38	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
140. S.×kapsheviakii_s40	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
141. S.×kapsnosperia_sc20a_NC	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
142. S.×kapsnosperia_sh2_NC	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
143. S.ochroleuca_4n_NY	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
144. S.ochroleuca_4wx_OH	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
145. S.ochroleuca_16b_VA	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
146. S.ochroleuca_16c_NH	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
147. S.ochroleuca_16f_NH	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
148. S.ochroleuca_16g_MI	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
149. S.ochroleuca_16h_NY	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
150. S.arcisepala_4u_OH	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
151. S.arcisepala_sc30a_OH	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
152. S.arcisepala_4y_NY	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
153. S.arcisepala_4z_OH	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC
154. S.arcisepala_NY1_NY	TTCTTTTG	CTTC	TCCTTTTCATTCTTCA	GCTCAGGCGCC	GGGAAGATCACGGCTGCTGCTGGGTTGTAGAAGGAAGCGATGGACATGCGGTTGC

**Figure 2.** Examples of *ACO* gene sequence concatenations for selected *Spiranthes*. Samples 135–140 labeled “S. ×kapsheviakii” represent *a priori* interior lowland *S. sheviakii* from Hough and Young (2021), all other samples are from Pace and Cameron (2017). Samples 141 & 142 represent Appalachian *S. ×kapsnosperia*. Samples of *S. cernua* s.s. are included immediately above the highlighted box and samples of *S. ochroleuca* are included immediately below the highlighted box **A** examples of ambiguities in *a priori* *S. sheviakii* that correspond to nucleotide differences between *S. cernua* and *S. ochroleuca* (e.g., G, R, A) **B** examples of nucleotide states that are shared with *S. ochroleuca* but not *S. cernua* (e.g., left-most highlighted A & G), and additional examples of ambiguous states in *a priori* *S. sheviakii* that correspond to nucleotide differences between *S. cernua* and *S. ochroleuca* (e.g., G, R, and A).

*S. sheviakii*. Additionally, the *ACO* nucleotide ambiguity patterns for *S. sheviakii* are distinct from those of regionally sympatric *S. incurva* (Jenn.) M.C. Pace, *S. magnicamporum* Sheviak, or any other member of the *S. cernua* species complex (Fig. 2), indicating these species are not involved in the evolution of *S. sheviakii*. The nucleo-



tide ambiguity and nuclear/chloroplast discordant patterns are consistent across all samples of *S. ×kapnosperia* and *S. sheviakii* included in Pace and Cameron (2017) and Hough and Young (2021). Thus, *S. ×kapnosperia* and *S. sheviakii* share the same ancestral hybrid parentage of *S. cernua* × *S. ochroleuca*, but this parentage is expressed differently within the genomes of the two resulting regionally distinct hybrid populations (Table 2). Nomenclaturally, per The Code (Article H.4.1):

When all the parent taxa can be postulated or are known, a nothotaxon is circumscribed so as to include all individuals recognizably derived from the crossing of representatives of the stated parent taxa (i.e. not only the F1 but subsequent filial generations and also back-crosses and combinations of these). There can thus be only one correct name corresponding to a particular hybrid formula; this is the earliest legitimate name (Art. 6.5) at the appropriate rank (Art. H.5), and other names corresponding to the same hybrid formula are synonyms of it.

Thus, any recognizably intermediate individual or population that results from the hybridization of *S. cernua* and *S. ochroleuca* must be recognized by the priority name *S. ×kapnosperia*, even if different hybridization events between the parental species occurred at different geologic times, in different places, resulting in different genomic expressions, and different morphologies. Based on the available *ndhJ* and *ACO* molecular data of Hough and Young (2021), *S. sheviakii* is unambiguously of hybrid origin between *S. cernua* and *S. ochroleuca*, and is thus synonymous with *S. ×kapnosperia*. It should be noted that species and nothospecies are the same nomenclatural rank, and the use of the multiplication symbol (×) is simply to emphasize the hybrid origin of nothospecies. Because Hough and Young (2021) appear to have been aware of the full hybrid parentage of their newly proposed name when they described *S. sheviakii* (based on the label of the type specimens, *Hough s.n.*, Fig. 1), this name is likely superfluous, although it is not illegitimate as they did not include the type of *S. ×kapnosperia* within the circumscription of *S. sheviakii*.

The evolutionary history of *S. ×kapnosperia* in its newly expanded understanding (*S. ×kapnosperia* sensu nov.) is perhaps one of the more unusual within the entire genus, having formed from the same two parental species (at least) two times, in widely disjunct locations, displaying different molecular signals between the parents. Additionally, the parental species likely played different maternal vs. paternal roles in the formation of the regionally disjunct *S. ×kapnosperia* populations (Table 2). The resulting differences in ambiguity patterns (or the lack of ambiguities) are likely due to differences in the hybridization and introgression histories of these regional populations. As Appalachian *S. ×kapnosperia* lacks *ACO* and nrITS ambiguities, it may be the result of chloroplast capture. This is a process through which an initial F1 hybridization event between paternal *S. cernua* and maternal *S. ochroleuca* is then followed by several backcrossing events with *S. cernua* as the pollen (paternal) parent, until the entire nuclear genome is only represented by *S. cernua*, but the chloroplast genome retains the original chloroplast contribution of *S. ochroleuca*. By contrast, the ambiguities present in the *ACO* locus of Interior Lowland *S. ×kapnosperia* (previously referred to as *S. sheviakii*) indicate it likely resulted from an initial F1 hybridization without exten-



sive (or only limited) backcrossing. Elsewhere in *Spiranthes*, Arft and Ranker (1998) hypothesized that at least two separate hybridization events between *S. magnicamporum* and *S. romanzoffiana* Cham. led to the formation of *S. diluvialis*, with subsequent localized dispersal in Utah and Colorado. However, the examined molecular signals from all sampled populations were the same (at the time of their study *S. diluvialis* was known from Colorado, Montana, Nevada, Utah, and Wyoming, but their study focused on samples from Colorado and Utah; Arft and Ranker 1998). Additional molecular phylogenetic study has not found major molecular differentiation between different populations of *S. diluvialis* (Dueck et al. 2014; Pace 2015).

*Spiranthes*  $\times$  *kapnosperia* was originally known to occur diffusely over a small region of the greater Smoky Mountain region and southern Blue Ridge Mountains, in the southern Appalachian Mountains of North Carolina, South Carolina, and Tennessee (Pace and Cameron 2017). The expanded understanding of *S.*  $\times$  *kapnosperia* sensu nov. discussed here extends the known distribution of this nothospecies throughout the distributional contact zone between *S. cernua* and *S. ochroleuca* along the northern limit of *S. cernua* in the area of the Interior Lowlands, Ohio River Valley, and southern Great Lakes Basin, an area that was not heavily sampled in the molecular work of Pace and Cameron (2017). Ecologically, populations in the southern Appalachians occur in more mesic sites vs. less mesic habitat of the Interior Lowlands populations; habitat variability is not uncommon across the genus. Morphologically, both disjunct populations are readily identifiable as intermediate hybrids of *S. cernua*  $\times$  *S. ochroleuca*. However, they display slightly different morphological affinities to their parents, with southern Appalachian *S.*  $\times$  *kapnosperia* being more similar to *S. ochroleuca*, and Interior Lowland *S.*  $\times$  *kapnosperia* being more similar to *S. cernua*. The flowers of southern Appalachian *S.*  $\times$  *kapnosperia* are generally slightly ascending (as is common in *S. ochroleuca*). The flowers of Interior Lowlands *S.*  $\times$  *kapnosperia* are generally very similar in overall size and appearance to *S. cernua* s.s., commonly with a nod to the flowers, but sharing the yellowish labellum coloration and rounded abaxial labellum glands with *S. ochroleuca* (Table 1).

Within the *S. cernua* species complex, molecular data have supported hybridization as a strong driver of speciation, with four of the seven non-hybrid species within the complex involved in the evolution of six species of hybrid origin or nothospecies (Table 3, Pace and Cameron 2017). *Spiranthes cernua* is the most frequently involved species, giving rise to the evolution of four hybrid taxa, and is typically the inferred maternal parent. The frequent involvement of *S. cernua* in the evolution of hybrid taxa may be due to its broad geographic distribution, stretching from Maritime Canada south to northern Florida and west through the mid- and southern-Appalachian Mountains to Texas. The repeated evolution of hybrid taxa such as *S.*  $\times$  *kapnosperia*, in addition to the cryptic morphological nature of the species within the complex, has contributed to the systematic and nomenclatural challenges commonly associated with the genus. The repeated evolution of *S.*  $\times$  *kapnosperia* and complicated hybridization history of the wider *S. cernua* species complex also highlight the need for slow and careful study when deciding to name and describe new taxa within the genus (Ames



**Table 3.** Ancestry of known hybrid taxa derived from members of the *S. cernua* species complex, as supported by combined molecular and morphological evidence.

Hybrid taxa	Inferred paternal species	Inferred maternal species	Literature source
<i>S. bightensis</i>	<i>S. odorata</i>	<i>S. cernua</i>	Pace (2021)
<i>S. casei</i>	<i>S. lacera</i> (var. <i>lacera</i> )	<i>S. ochroleuca</i>	Pace (2015); Pace unpublished data
<i>S. diluvialis</i>	<i>S. magnicamporum</i>	<i>S. romanzoffiana</i>	Arft and Ranker (1998)
<i>S. incurva</i>	<i>S. magnicamporum</i>	<i>S. cernua</i>	Pace and Cameron (2017)
<i>S. ×kapnosperia</i> sensu nov.	<i>S. cernua</i> (southern Appalachian populations) or <i>S. ochroleuca</i> (Interior Lowland populations)	<i>S. cernua</i> (Interior Lowland populations) or <i>S. ochroleuca</i> (southern Appalachian populations)	Pace and Cameron (2017); Hough and Young (2021)
<i>S. niklasii</i>	<i>S. cernua</i>	<i>S. ovalis</i>	Pace and Cameron (2017)


1921), with particular attention given to nomenclatural rules and priority. Although they do not provide any molecular data, Hough and Young (2021), using terminology from Sheviak (1982), also place emphasis on the potential distinction of “low prairie race” and “southern prairie complex” populations currently contained within the circumscription of *S. cernua* s.s. Moving forward, researchers should keep in mind the priority names for taxa that involve hybridization between *S. cernua* and other members of the *S. cernua* species complex (Table 3). Additionally, not all individuals or populations of hybrid ancestry should be named, as genomic data increasingly shows complex hybridization and introgression patterns make for cryptically complex genetic ancestries and species relationships in groups with porous reproductive barriers (e.g., Evans et al. 2023).

A few additional notes related to Hough and Young (2021) are discussed here: 1) Hough and Young (2021) discuss the “holotype for *S. incurva*.” As detailed in Pace and Cameron (2017), the name *S. incurva* is a nomenclatural combination based on the basionym *Ibidium incurvum* Jenn.:

Since Jennings selected a suite of specimens, “Aug. 24–26, 1905”, housed at CM as “the type specimens”, and not a specific specimen, collection number, or sheet, the specimen designated by Catling as the holotype, via an annotation label, is more properly designated as the lectotype. All other specimens collected on Aug. 26, 1905 must then be isolectotypes, and all other specimens collected within “the type specimens” collection range designated as syntypes.

The lectotype of *I. incurvum* is the *Jennings s.n.* specimen collected on 26 August 1905, from Fog Whistle (CM). However, this specimen is not discussed or examined in Hough and Young (2021), which only discusses the remaining syntypes. 2) Hough and Young (2021) note “at least within the range of this study, we have not observed *S. incurva* growing in xeric sites. The typical habitats appear to be mostly moist to wet and mediacid to calcareous.” It should be noted that *S. incurva* is found in a wide variety of habitats, from hot, dry, sandy lake beach dunes, old fields, and roadside embankments, to standing in shallow water of fens and lake beach dune swales (Pace and Cameron 2017). This is inclusive of locations within the study range of Hough and Young (2021). 3) Hough and Young (2021) make much of apparent ambiguities





**Figure 3.** Examples of *ACO* gene nucleotide ambiguities for selected *Spiranthes*. The sample “*S. arcisepala\_S4*” is from Hough and Young (2021), all other samples are from Pace and Cameron (2017) **A** examples of ambiguities present in Hough and Young (2021), but not present in Pace and Cameron (2017) (Y and R), and an ambiguity present in Hough and Young (2021) but overlooked in Pace and Cameron (2017) (K) **B** example of ambiguity present in Pace and Cameron (2017) (M), but not present in Hough and Young (2021).

in the nuclear *ACO* data for their accessions of *S. arcisepala* M.C. Pace, claiming that Pace and Cameron (2017) misinterpreted their data. After comparing the GenBank data of Hough and Young (2021) to the raw sequence data of Pace and Cameron (2017) (Fig. 3), there was a single instance where Pace and Cameron (2017) missed an ambiguity that is present in the data of Hough and Young (2021). However, the overwhelming majority of supposed *S. arcisepala* ambiguities present in Hough and Young (2021) are simply not present in our data, which are unambiguously a single nucleotide. Furthermore, I found an additional ambiguity in the *ACO* data of Pace and Cameron (2017) that is not present in the *ACO* data of Hough and Young (2021), and this point of ambiguity does not correspond to a point of molecular differentiation between any other member of the *S. cernua* species complex (Fig. 3). Based on this reexamination and comparison, I reassert that *S. arcisepala* is not of hybrid origin, although it may be an autopolyploid. Genomic examination of hybridization across the genus is obviously needed, and is currently underway.



## Nomenclature

***Spiranthes* ×*kapnosperia* M.C. Pace [*S. cernua* × *S. ochroleuca*], Syst. Bot. 42: 659. 2017.**

= *Spiranthes sheviakii* M. Hough and M.A. Young. Native Orchid Conf. J. 18.3: 35. 2021. Type: U.S.A. New York: Onondaga County, town of Lysander, Three Rivers WNA, 19 Sep 2020, *M. Hough and M.A. Young s.n.* (holotype: BH! [BH 000298396]; isotype: CORT! (×2)).

**Type.** U.S.A. North Carolina: Transylvania County, Great Smoky Mountains, Pisgah National Forest, ca 7.5 km NW of Balsam Grove, north side of 215, below a steep seeping cliff, growing in moss and lichen hummocks, 2 Oct 2016, *M.C. Pace 1030* (holotype: NY! [01392730]; isotype: NCU! [NCU00332163], US!).

## Acknowledgements

Thank you to Kanchi Gandhi (Harvard University) for helpful conversations regarding the nomenclature of hybrids while this manuscript was in preparation; two anonymous reviewers for their helpful comments; Anna M. Stalter (BH) and Michael Hough (CORT) for sending images of the *S. sheviakii* types. James (Jim) Fowler (1947–2021) provided samples that led to the description of *S. ×kapnosperia*.

## References

- Ackerman JD, Galarza-Pérez M (1991) Patterns and maintenance of extraordinary variation in the Caribbean orchid, *Tolumnia* (*Oncidium*) *variegata*. Systematic Botany 16(1): 182–194. <https://doi.org/10.2307/2418982>
- Ackerman JD, Phillips RD, Tremblay RL, Karremans A, Reiter N, Peter CI, Bogarín D, Pérez-Escobar OA, Liu H (2023) Beyond the various contrivances by which orchids are pollinated: global patterns in orchid pollination biology. Botanical Journal of the Linnean Society XX: 1–30. <https://doi.org/10.1093/botlinnean/boac082>
- Ames O (1921) Notes on New England orchids. I. *Spiranthes*. Rhodora 23: 3–85.
- Arft AM, Ranker TA (1998) Allopolyploid origin and population genetics of the rare orchid *Spiranthes diluvialis*. American Journal of Botany 85(1): 110–122. <https://doi.org/10.2307/2446560>
- Arnold ML (2001) Anderson's paradigm: Louisiana Irises and the study of evolutionary phenomena. Molecular Ecology 9(11): 1687–1698. <https://doi.org/10.1046/j.1365-294x.2000.01090.x>
- Dueck LA, Aygoren D, Cameron KM (2014) A molecular framework for understanding the phylogeny of *Spiranthes* (Orchidaceae), a cosmopolitan genus with a North American center of diversity. American Journal of Botany 101(9): 1551–1571. <https://doi.org/10.3732/ajb.1400225>



- Evans SA, Whigham DF, Hartvig I, McCormick MK (2023) Hybridization in the Fringed Orchids: An Analysis of Species Boundaries in The Face of Gene Flow. *Diversity* (Basel) 15(3): 384. <https://doi.org/10.3390/d15030384>
- Fu PC, Sun SS, Khan G, Dong XX, Tan JZ, Favre A, Zhang FA, Chen ZL (2020) Population subdivision and hybridization in a species complex of *Gentiana* in the Qinghai-Tibetan Plateau. *Annals of Botany* 125(4): 677–690. <https://doi.org/10.1093/aob/mcaa003>
- Gleason HA, Cronquist A (1963) Manual of vascular plants of Northeastern United States and adjacent Canada. D. Van Nostrand Company.
- Hough M, Young MA (2021) A systematic survey of the *Spiranthes cernua* species complex (Orchidaceae) in New York. *The Native Orchid Conference Journal* 18(3): 22–56.
- Jacquemyn H, Brys R, Honnay O, Roldán-Ruiz I, Lievens B, Wiegand T (2012) Nonrandom spatial structuring of orchids in a hybrid zone of three *Orchis* species. *The New Phytologist* 193(2): 454–464. <https://doi.org/10.1111/j.1469-8137.2011.03913.x>
- Pace MC (2015) Evolution, species boundaries, and biogeography of *Spiranthes* (Orchidaceae): Uncoiling Ames’s “most perplexing orchid genus in our flora”. Ph.D. dissertation. University of Wisconsin, Madison, WI.
- Pace MC (2021) *Spiranthes bightensis* (Orchidaceae), a New and Rare Cryptic Hybrid Species Endemic to the U. S. Mid-Atlantic Coast. *Phytotaxa* 498(3): 159–176. <https://doi.org/10.11646/phytotaxa.498.3.2>
- Pace MC, Cameron KM (2017) The systematics of the *Spiranthes cernua* species complex (Orchidaceae): Untangling The Gordian Knot. *Systematic Botany* 42(4): 640–669. <https://doi.org/10.1600/036364417X696537>
- Pace MC, Cameron KM (2019) The evolutionary and systematic significance of hybridization between taxa of *Spiranthes* (Orchidaceae) in the California Sierra Nevada and Cascade Range. *Taxon* 68(2): 199–217. <https://doi.org/10.1002/tax.12034>
- Pace MC, Freudenstein JV (2018) Orchidaceae, the orchid family. In: Naczi RFC, et al. (Eds) New Manual of Vascular Plants of Northeastern United States and Adjacent Canada, online edition 2016 onward. NYBG Press, New York. <https://doi.org/10.21135/893275471.096>
- Pace MC, Giraldo G, Frericks J, Lehnebach CA, Cameron KM (2019) Illuminating the systematics of the *Spiranthes sinensis* species complex (Orchidaceae): Ecological speciation with little morphological differentiation. *Botanical Journal of the Linnean Society* 189(1): 36–62. <https://doi.org/10.1093/botlinnean/boy072>
- Pillon Y, Fay MF, Hedrén M, Bateman RM, Devey DS, Shipunov AB, van der Bank M, Chase MW (2007) Evolution and temporal diversification of western European polyploid species complexes in *Dactylorhiza* (Orchidaceae). *Taxon* 56(4): 1185–1208. <https://doi.org/10.2307/25065911>
- Pinheiro F, De Barros F, Palma-Silva C, Meyer D, Fay MF, Suzuki RM, Lexer C, Cozzolino S (2010) Hybridization and introgression across different ploidy levels in the Neotropical orchids *Epidendrum fulgens* and *E. puniceoluteum* (Orchidaceae). *Molecular Ecology* 19(18): 3981–3994. <https://doi.org/10.1111/j.1365-294X.2010.04780.x>
- Sheviak CJ (1982) Biosystematic study of the *Spiranthes cernua* complex. Bulletin of the New York State Museum Science Service, Bulletin Number 448. <https://doi.org/10.5962/bhl.title.135544>



- Sheviak CJ (1991) Morphological variation in the compliospecies *Spiranthes cernua* (L.) Rich.: Ecologically-limited effects of gene flow. *Lindleyana* 6: 228–234.
- Sheviak CJ, Brown PM (2002) *Spiranthes*. In: Flora of North America Editorial Committee (Eds) Flora of North America vol. 26. Oxford University Press, New York, 530–545.
- Sheviak CJ, Catling PM (1980) The identity and status of *Spiranthes ochroleuca* (Rydberg) Rydberg. *Rhodora* 82: 525–562.
- Soliva M, Widmer A (2003) Gene flow across species boundaries in sympatric, sexually deceptive *Ophrys* (Orchidaceae) species. *Evolution; International Journal of Organic Evolution* 57(10): 2252–2261. <https://doi.org/10.1111/j.0014-3820.2003.tb00237.x>
- Sun M (1996) The allopolyploid origin of *Spiranthes hongkongensis* (Orchidaceae). *American Journal of Botany* 83(2): 252–260. <https://doi.org/10.1002/j.1537-2197.1996.tb12702.x>
- Surveswaran S, Gowda V, Sun M (2018) Using an integrated approach to identify cryptic species, divergence patterns and hybrid species in Asian ladies' tresses orchids (*Spiranthes*, Orchidaceae). *Molecular Phylogenetics and Evolution* 124: 106–121. <https://doi.org/10.1016/j.ympev.2018.02.025>
- Szalanski AL, Steinauer G, Bischof R, Petersen J (2001) Origin and conservation genetics of the threatened Ute ladies'-tresses, *Spiranthes diluvialis* (Orchidaceae). *American Journal of Botany* 8(1): 177–180. <https://doi.org/10.2307/2657138>
- Wettewa E, Bailey N, Wallace L (2020) Comparative Analysis of Genetic and Morphological Variation within the *Platanthera hyperborea* Complex (Orchidaceae). *Systematic Botany* 45(4): 767–778. <https://doi.org/10.1600/036364420X16033962925303>